

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
7 April 2005 (07.04.2005)

PCT

(10) International Publication Number
WO 2005/030969 A1

(51) International Patent Classification⁷: **C12N 15/85**

(21) International Application Number:
PCT/KR2004/002474

(22) International Filing Date:
24 September 2004 (24.09.2004)

(25) Filing Language: Korean

(26) Publication Language: English

(30) Priority Data:
10-2003-0068641 2 October 2003 (02.10.2003) KR

(71) Applicant (for all designated States except US): **PROGEN CO., LTD.** [KR/KR]; 47-5 Sanggal-ri Giheung-eup Yon-gin-si, Kyunggi-do 449-900 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YANG, Se Hwan** [KR/KR]; #5-305 Nakwon Apt. Jigok-dong Nam-ku Pohang-si, Gyeongsangbuk-do 790-751 (KR). **SUNG, Young Chul** [KR/KR]; #101-601 Hyundai Edong Hometown Daejam-dong Nam-ku Pohang-si, Gyeongsangbuk-do

790-310 (KR). **NA, Kyu-Heum** [KR/KR]; #312-1303 Cheongmyeong Dongshin Apt. Yeongtong-dong Paldal-ku Suwon-si, Kyunggi-do 442-470 (KR). **LEE, Sung Hee** [KR/KR]; #101-306 Nakseongdae Hyundai Apt. 179-58 Bongcheon-11dong Gwanak-ku, Seoul 151-815 (KR). **KIM, Won Bae** [KR/KR]; #1801 Daelim Acrovill Apt. 467-6 Dogok-dong Kangnam-ku, Seoul 135-270 (KR).

(74) Agent: **LEE, Won-Hee**; 8th Fl. Sung-ji Heights II 642-16 Yoksam-dong Kangnam-ku, Seoul 135-080 (KR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: METHOD FOR MASS PRODUCTION OF HUMAN FOLLICLE STIMULATING HORMONE

Wild type DNA sequence	ATG AAG ACA CTC CAG TTT TTC TTC CTT TTC TGT TGC TGG AAA GCA
Amplified DNA sequence	---
Amino acid sequence	Met Lys Thr Leu Gln Phe Phe Phe Leu Phe Cys Cys Trp Lys Ala
	ATC TGC TGC AAT AGC TGT GAG CTG ACC AAC ATC ACC ATT GCA ATA
	Ile Cys Cys Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile
	GAG AAA GAA GAA TGT CGT TTC TGC ATA AGC ATC AAC ACC ACT TGG
	Glu Lys Glu Glu Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp
	TGT GCT GGC TAC TGC TAC ACC AGG GAT CTG GTG TAT AAG GAC CCA
	Cys Ala Gly Tyr Cys Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro
FSH β subunit	
	GCC AGG CCC AAA ATC CAG AAA ACA TGT ACC TTC AAG GAA CTG GTA
	Ala Arg Pro Lys Ile Gln Lys Thr Cys Thr Phe Lys Glu Leu Val
	TAT GAA ACA GTG AGA GTG CCC GGC TGT GCT CAC CAT GCA GAT TCC
	Tyr Glu Thr Val Arg Val Pro Gly Cys Ala His His Ala Asp Ser
	TTG TAT ACA TAC CCA GTG GCC ACC CAG GTG CAC TGT GGC AAG TGT
	Leu Tyr Thr Tyr Pro Val Ala Thr Gln Val His Cys Gly Lys Cys
	GAC AGC GAC AGC ACT GAT TGT ACT GTG CGA GGC CTG GGG CCC AGC
	Asp Ser Asp Ser Thr Asp Cys Thr Val Arg Gly Leu Gly Pro Ser
	TAC TGC TCC TTT GGT GAA ATG AAA GAA TAA
	Tyr Cys Ser Phe Gly Glu Met Lys Glu termination

(57) Abstract: The present invention relates to a method for mass-production of human follicle stimulating hormone, more precisely, to an expression vector containing human follicle stimulating hormone gene, a transformant transfected with the expression vector and a method for mass-production of human follicle stimulating hormone by using the same. Human follicle stimulating hormone can be produced largely by using an expression vector and a transformant of the present invention. Therefore the expression vector and the transformant of the present invention can be effectively used for the treatment of infertility.

WO 2005/030969 A1



European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

— *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

METHOD FOR MASS PRODUCTION OF HUMAN
FOLLICLE STIMULATING HORMONE

5

Technical Field

The present invention relates to a method for mass-production of human follicle stimulating hormone, more precisely, to an expression vector containing human follicle stimulating hormone gene, a transformant
10 transfected with the expression vector and a method for mass-production of human follicle stimulating hormone by using the same.

Background Art

15

Follicle stimulating hormone (referred as "FSH" hereinafter), having a molecular weight of 32,600 Da, is a glycoprotein secreted in anterior pituitary, more precisely, a heterodimeric glycoprotein in which alpha and beta subunits are combined each other by a non-
20 covalent bond. Heterodimeric glycoproteins are exemplified by FSH, luteinizing hormone (referred as "LH" hereinafter) which is a pituitary glycoprotein, human chorionic gonadotropin (referred as "hCG" hereinafter) which is a placental glycoprotein, thyroid

stimulating hormone (referred as "TSH" hereinafter), factor VIII and IL-12, etc. The alpha subunit of FSH is very similar in structure and in immunological aspects to LH, TSH and hCG. But, the beta subunit is different from every hormone, meaning that each hormone has unique biological and immunological characteristics owing to the different properties of beta subunit (Pierce. J.G and Parsons. T.F., *Ann. Rev. Biochem.*, 50:465-495, 1981). In particular, human FSH (hFSH) stimulates follicle of an ovary to make the follicle be growing, leading to the ovulation. Thus, it has been in clinical use for the treatment of sterility. FSH is obtained from urine of pregnant women, but the obtainable amount is very small and the separation is difficult. Therefore, it is urgently required to develop a method for mass-production of the FSH by using a genetic recombinant technique.

In order to prepare a transformed cell line expressing a glycoprotein hormone by taking advantage of a genetic recombinant technique, *E. coli*, yeast, insect cells or animal cells can be used as a host cell. When the selection of a host cell is made to prepare a glycoprotein, glycosylation capacity or glycosylation system must be considered first because every host cell has a different glycosylation capacity or glycosylation

system. Especially to prepare a human FSH, it is required to select a host cell with an excellent glycosylation capacity because glucose plays an important role in *in vivo* activity of hormone. For example, among the above-mentioned candidates for a host cell, *E. coli* has no glycosylation capacity at all. Although yeast or insect cells have a little glycosylation capacity, it is too low to produce a complex type glycoprotein such as human FSH, or it can only produce high mannose type glycoprotein. Therefore, they are not promising candidates for a host cell. Even though mammalian cells have a very good glycosylation capacity or glycosylation system, the level or the type of glycosylation varies from the characteristics of each cell line. So, in order to produce a glycoprotein such as hFSH available for the treatment of a disease, such a mammalian cell line that does not cause antigenicity but is safe and has high *in vivo* activity must be selected as a host cell.

It is also essential to develop an efficient expression vector for the mass-production of a heterodimeric glycoprotein such as FSH, LH, hCG, TSH, factor VIII and IL-12 by using animal cells. These are the examples of a constituent of such expression vector; promoter playing an important role in

controlling the gene expression, intron enabling the
increase of transcription efficiency of a gene by
stabilizing mRNA, polyadenylation motif (pA) enabling
the increase of the stability of mRNA, and selection
5 marker, etc.

Thus, the present inventors prepared an
expression vector highly and stably expressing human
FSH, and then completed this invention by confirming
10 that FSH could be mass-produced stably by transfecting
a host cell having an excellent glycosylation capacity
with the prepared expression vector.

Disclosure

15 Technical Problem

It is an object of the present invention to
provide an expression vector containing human follicle
stimulating hormone gene, a transformant transfected
with the expression vector and a method for mass-
20 production of human follicle stimulating hormone by
using the same.

Technical Solution

In order to achieve the above object, the present

invention provides a recombinant expression vector containing human FSH gene.

The present invention also provides a transformant transfected with the expression vector.

5 The present invention further provides a method for mass-production of human follicle stimulating hormone by using the transformant.

10 Hereinafter, the present invention is described in detail.

The present invention provides a recombinant expression vector containing human FSH gene.

15 In order to mass-produce heterodimeric glycoprotein such as FSH, LH, hCG, TSH, factor VIII and IL-12 using animal cells, it is essential to develop an efficient expression vector. Thus, the present invention provides a highly efficient expression vector for mass-production of human FSH, considering all the factors involved.

20

1) A gene coding hFSH is composed of two subunits (alpha and beta subunits) which are transcribed from different genes. In the preferred embodiment of the present invention, IRES (internal ribosomal entry site)
25 of EMCV (encephalomyocarditis virus) was used in order

to prepare a heterodimeric glycoprotein by expressing two different glycoproteins in the same cell. Then the resultant was inserted in between alpha subunit gene and beta subunit gene of hFSH to make those genes be expressed simultaneously. So, it is preferred for a gene coding hFSH included in the expression vector of the present invention to contain FSH alpha subunit gene represented by SEQ. ID. No 1 and FSH beta subunit gene represented by SEQ. ID. No 2, and it is more preferred for the gene coding hFSH to have IRES sequence, represented by SEQ. ID. No 7, inserted in between FSH alpha subunit gene and FSH beta subunit gene in order to induce simultaneous expression of both subunit genes.

2) A promoter plays an important role in the regulation of gene expression. So, an expression vector of the present invention was prepared to contain a promoter gene. A promoter included in the expression vector of the present invention is preferably the promoter of early gene of cytomegalovirus (CMV) represented by SEQ. ID. No 8 which has been known as the most powerful promoter.

3) Mammalian genes contain intron. The expression level is remarkably increased in the presence of intron (Korb *et al.*, *Nucleic Acids Res*, 25:5901-5908, 1993). This might be because that intron

can stabilize mRNA and increase transcription efficiency. Thus, an expression vector of the present invention was designed to have intron sequence, and the intron sequence, at that time, was preferably tripartite leader sequence (referred as "TPL" hereinafter) of adenovirus represented by SEQ. ID. No 9.

4) In order to increase the stability of mRNA, polyadenylation motif (pA) is important (Drummond et al., *Nucleic Acids Res*, 25:7375-7394, 1985). The expression vector of the present invention includes such polyadenylation motif, and the polyadenylation motif is preferred to be polyadenylation motif of early gene of SV40 virus, represented by SEQ. ID. No 13, and/or polyadenylation motif of bovine growth hormone (BGH) gene, represented by SEQ. ID. No 14. In the preferred embodiment of the present invention, an expression vector was prepared by locating BGH pA behind FSH alpha gene, SV40 pA behind Neo gene, and SV40 pA gene behind dihydrofolate reductase (referred as "DHFR") gene (see FIG. 1).

5) A selection marker gene is necessary to select cell lines that can over-express FSH after being transfected with the expression vector of the present invention. Thus, the expression vector of the present invention was designed to include a selection marker

gene, and for the selection marker, DHFR gene, a selection marker represented by SEQ. ID. No 12, was preferably used to facilitate selection of a cell line from CHO/dhfr⁻ cells which were Chinese hamster ovary (CHO) originated cell line. CHO/dhfr⁻ cell line has damaged DHFR gene, indicating that its' nucleic acid synthesizing process is incomplete. But, when the cell is transfected with FSH expression vector of the present invention containing DHFR gene, nucleic acid synthesis can be completed. Thus, it is possible to select those cells expressing FSH by the introduction of FSH expression vector.

Taking all the above-mentioned factors into consideration, the present inventors prepared a recombinant expression vector to express human FSH that includes 1) a gene coding human FSH, 2) a promoter sequence, 3) an intron sequence, 4) a polyadenylation motif sequence, and 5) a selection marker gene.

In the preferred embodiment of the present invention, in order to express both FSH alpha and beta subunits simultaneously in the same cell, IRES was inserted in between FSH alpha subunit gene and FSH beta subunit gene. And the expression efficiency of hFSH was maximized by using a promoter of early gene of cytomegalovirus that has been known as the most

powerful promoter. Also, adenovirus tripartite leader sequence (TPL) containing intron to increase the stability and transcription efficiency and polyadenylation motif of early gene of SV40 virus as well as polyadenylation motif of bovine growth hormone gene were all included in the vector of the present invention. In addition, DHFR gene was included as a selection marker to facilitate the selection of cell lines over-expressing FSH from the cells transfected with the expression vector of the present invention.

The present inventors prepared an expression vector containing all the above-mentioned elements and named it 'Rc/CMV-dhfr-TPL-hFSH beta/alpha'.

The expression vector of the present invention was prepared for the high-expression of human FSH gene, a heterodimeric glycoprotein. And the vector of the invention is very efficient one enabling mass-production of a heterodimeric glycoprotein. It is well known to the people in this field that instead of human FSH gene, any other heterodimeric glycoprotein such as LH, hCG, TSH, factor VIII and IL-12 gene can be included in the expression vector and the resultant vector can be effectively used for the mass-production of a protein coded from the above genes.

25

The present invention also provides a transformant transfected with the expression vector.

In order to prepare a cell line high-expressing human FSH, which is highly activated in a human body, the present inventors selected Chinese hamster ovary (referred as "CHO" hereinafter) cells among many mammalian cell lines as a host cell line because they have a satisfactory glycosylation capacity or glycosylation system, do not induce antigenicity *in vivo*, are safe and have a carbohydrate structure showing a high *in vivo* activity.

Nucleic acid biosynthesizing processes occurring in CHO/dhfr⁻ cells, originated from CHO, are incomplete because DHFR gene therein is damaged. But, nucleic acid biosynthesis is fully recovered in CHO cells when they are transfected with the FSH expression vector of the present invention (Rc/CMV-dhfr-TPL-hFSH beta/alpha) that contains DHFR gene. Therefore, it is possible to select those cells harboring the expression vector by culturing CHO cells in a cell culture medium devoid of nucleic acid. And the cells are growing in a culture medium supplemented with methotrexate (MTX), a substrate analogue of DHFR, by the amplification of DHFR gene in cells. At this time, the amplification of FSH gene is also performed because FSH expression

vector of the present invention contains DHFR gene. Thus, adding MTX, a substrate analogue of DHFR, into a cell culture medium can improve the expression level of FSH protein in FSH expressing cells.

5 In the present invention, CHO/dhfr⁻ cells were transfected with FSH expressing vector, and clones which were highly expressed in high-concentration of MTX and showed homologous were selected and further cultured, resulting in a cell line in which FSH was
10 stably and highly expressed (see FIG. 4 to FIG. 6). The selected cell line high-expressing FSH was named "DPFC325" which was deposited at Korean Cell Line Research Foundation (KCLRF) on June 20, 2003 (Accession No: KCLRF-BP-00082).

15

The present invention also provides a method for mass-production of human follicle stimulating hormone by using the transformant.

20 The method for mass-production of human FSH of the present invention includes the following steps:

- 1) Preparation of an expression vector containing human FSH gene;
- 2) Transfection of host cells with the expression vector of the step 1);
- 25 3) Selection of transformants transfected in the

step 2);

4) Selection of a recombinant transformant stably producing FSH from the recombinant transformants selected in the step 3); and

5) Obtainment of FSH from the culture of the recombinant transformant selected in the step 4).

In step 1), the expression vector preferably includes human FSH coding gene, a promoter sequence, an intron sequence, a polyadenylation motif sequence and a selection marker gene, and is more preferably "Rc/CMV-dhfr-TPL-hFSH beta/alpha" constructed in the present invention.

In step 2), the host cell is preferably CHO/dhfr⁻ cell, a CHO originated cell line having incomplete nucleic acid biosynthesizing processes because of the damaged DHFR gene.

In step 4), the recombinant transformant stably producing FSH is preferably "DPFC325" which was deposited at Korean Cell Line Research Foundation (KCLRF) on June 20, 2003 (Accession No: KCLRF-BP-00082).

In step 5), the obtainment of produced FSH is performed by conventional protein purification methods.

Description of Drawings

FIG. 1 is a schematic diagram showing the

preparation process of Rc/CMV-DHFR-TPL-hFSH beta/alpha
vector of the present invention,

FIG. 2 shows cDNA nucleotide sequence and amino
5 acid sequence of hFSH-alpha subunit expressed by pSK-
FSH-alpha vector of the present invention,

FIG. 3 shows cDNA nucleotide sequence and amino
acid sequence of hFSH-beta subunit expressed by pSK-
10 FSH-beta vector of the present invention,

FIG. 4 is a graph showing the expression of human
FSH after transfecting COS-7 cells with Rc/CMV-dhfr-
TPL-hFSH beta/alpha vector, confirmed by enzyme-linked
15 immunosorbent assay (ELISA),

FIG. 5 is a graph showing the result of enzyme-
linked immunosorbent assay (ELISA), confirming the
expression level of human FSH in clones that were
20 selected from cells adapted to 1 uM of MTX by limiting
dilution,

FIG. 6 is a photograph showing the result of FISH,
confirming whether or not human FSH gene is inserted in
25 chromosome of a CHO cell.

Mode for Invention

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

<Example 1> Construction of a recombinant hFSH expression vector (Rc/CMV-dhfr-TPL-hFSH beta/alpha)

<1-1> Preparation of human FSH alpha subunit and beta subunit

In order to prepare a structural gene that is essential for the production of a cell line expressing a recombinant human FSH, human FSH alpha subunit and beta subunit genes obtained from human pituitary gland cDNA library were amplified by PCR. More precisely, human pituitary gland cDNA library (cal#HL1139a, Clontech, USA) was used as a substrate, and primers were prepared for PCR with nucleotide sequence of FSH alpha subunit gene represented by SEQ. ID. No 1 and

nucleotide sequence of FSH beta subunit gene represented by SEQ. ID. No 2. The primer sequences for PCR amplification of FSH alpha and beta subunit genes were as follows.

5 1) FSH alpha subunit

Sense primer (SEQ. ID. No 3):
AGCGCCATGGATTACTACAGAAAATAT (underlined region : Nco I
recognition site),

Antisense primer (SEQ. ID. No 4):
10 GGGGGATCCGGGCCCTTAAGATTTGTGATAATTAACA (underlined
region : BamH I and Apa I recognition site)

2) FSH beta subunit

Sense primer (SEQ. ID. No 5):
GGGGGCGGCCGCAGGATGAAGACACTCCAGTTT (underlined region :
15 Not I recognition site),

Antisense primer (SEQ. ID. No 6):
GGGGATATCTTATTCTTTCATTTACACC (underlined region : EcoR
V recognition site)

20 The whole fraction of pituitary gland cDNA
library was used as a substrate, and each pair of
primers (alpha subunit: SEQ. ID. No 3 and No 4, beta
subunit: SEQ. ID. No 5 and No 6) were used for PCR.
PCR was performed as follows; predenaturation at 94°C
25 for 30 seconds, denaturation at 94°C for 30 seconds,

annealing at 57°C for 30 seconds, polymerization at 72°C for 30 seconds, 30 cycles from denaturation to polymerization, and final extension at 72°C for 5 minutes. The same PCR condition as described above was applied to the amplification of both FSH alpha subunit and FSH beta subunit.

<1-2> Construction of a vector for simultaneous expression of FSH alpha and beta subunit genes

FSH is composed of two subunits, which are FSH alpha subunit and FSH beta subunit. So, in order to construct an expression vector enabling simultaneous expression of both FSH alpha subunit and FSH beta subunit, IRES (internal ribosomal entry site) of EMCV (encephalomyocarditis virus), represented by SEQ. ID. No 7 was introduced, in which beta subunit was located at the upstream and alpha subunit was located at the downstream.

First, FSH beta subunit gene (390 nucleotides) which was amplified in the above example <1-1> was treated with T4 polynucleotide kinase, then cloned into pSK vector (Stratagene, USA), pretreated with Hinc II, by blunt ligation. The resultant vector was named "pSK-FSH-beta". Next, FSH alpha subunit gene (351

nucleotides), which had been amplified in the above example <1-1>, was digested with Nco I and BamH I, then inserted in the location of Nco I and BamH I of pSK-IRES vector (Ha et al., *Nat. Biotechnol.*, 20(4):381-386, 2002) containing EMCV IRES, resulting in "pSK-IRES-FSH-alpha". FSH-beta DNA fragment, prepared by digesting pSK-FSH beta vector with Xho I and EcoR V, was inserted in pSK-IRES-FSH-alpha vector, which had been digested with Xho I and EcoR V, resulting in the construction of a target vector containing both FSH-alpha subunit gene and FSH-beta subunit gene, which was named "pSK-FSH-alpha/IRES/FSH-beta".

Lastly, a vector was constructed to express FSH-alpha subunit gene and FSH-beta subunit gene simultaneously in animal cells. For the construction, Rc/CMV plasmid (Invitrogen, USA) harboring a very powerful cytomegalovirus (CMV) promoter, represented by SEQ. ID. No 8, was used as a basic vector. In order to increase the level of expression, adenovirus tripartite leader sequence (TPL), represented by SEQ. ID. No 9, was inserted in pTV2 vector (Lee et al., *J. Virol.*, 72(10):8430-8436, 1998). More precisely, pTV2 vector was used as a template and TPL gene was amplified by PCR using the below primers under the same PCR condition as described in the example <1-1>.

The primer sequences for the amplification of TPL gene are as follows.

Sense primer (SEQ. ID. No 10):
GATATCGATACTCTCTTCC,
5 Antisense primer (SEQ. ID. No 11):
GCGTCGACCTGCAGTTGGACCTGGGAG (underlined region : Sal I
recognition site)

The amplified TPL gene was treated with T4 polynucleotide kinase, which was then inserted in pSK
10 vector (Stratagene, USA) predigested with EcoR V. The
resultant product was digested with Sal I, so that the
location of Sal I in pSK vector which was corresponding
to the area of 5' of TPL gene and the location of Sal I
in 3' of TPL gene were restricted, resulting in the
15 separation of TPL gene. The separated gene was treated
with Klenow enzyme to make it have blunt ends. After
digesting with Hind III, Rc/CMV vector (Invitrogen, USA)
was also treated with Klenow enzyme to make it have
blunt ends, too. The treated vector was linked to the
20 separated TPL gene by the action of ligase, resulting
in RcCMV-TPL vector. For the amplification of the gene
expression, pSV2-dhfr vector (ATCC 37146) containing
mouse dihydrofolate reductase (DHFR) gene, represented
by SEQ. ID. No 12, and SV40 pA gene, represented by SEQ.
25 ID. No 13, in the rear of the above gene was digested

with Pvu II, then BamH I linker (SEQ. ID. No 15) was inserted in the position. The resultant product was digested with BamH I, resulting in 1.8 kb sized gene containing DHFR gene and SV40 pA gene. The gene was
5 inserted in RcCMV-TPL vector which had been digested with Bgl II, and the final vector was named "Rc/CMV-dhfr-TPL". PSK-FSH-alpha/IRES/FSH-beta vector was digested with Not I to prepare FSH-alpha/IRES/FSH-beta fragment. The fragment was inserted in the location of
10 Not I of Rc/CMV-dhfr-TPL vector to prepare an expression vector for high-expression of FSH. The product was named "Rc/CMV-dhfr-TPL-hFSH beta/alpha" (FIG. 1). DNA sequencing of FSH-alpha subunit gene and FSH-beta subunit gene in Rc/CMV-dhfr-TPL-hFSH
15 beta/alpha plasmid was performed by the method of Sanger et al. (Sanger et al., *Proc. Natl. Acad. Sci. USA*, 74:5463-5467, 1977).

As a result, DNA sequences of FSH-alpha subunit and FSH-beta subunit that were inserted in the
20 expression vector of the present invention were confirmed to be identical with the reported DNA sequences of FSH-alpha subunit and FSH-beta subunit, indicating that the expression vector of the present invention was correctly constructed (FIG. 2 and FIG. 3).

25

<Example 2> Preparation of a cell line expressing humanFSH

A cell line was transfected with the recombinant human FSH expression vector (Rc/CMV-dhfr-TPL-hFSH beta/alpha), constructed in the above example 1, to investigate the expression of recombinant human FSH. Particularly, COS-7 cells (ATCC, catalog # CRL-1651), being under culture, were inoculated into 60 mm dish plates (5×10^5 cells/plate), followed by further culture for 18 hours. The culture medium (DMEM + 10% FBS) was replaced with a fresh medium (4.5 ml). Then, about 4 hours later, the cells were transfected with a FSH expression vector Rc/CMV-dhfr-TPL-hFSH beta/alpha plasmid or a control vector Rc/CMV-dhfr-TPL plasmid by means of CaPO_4 coprecipitation. More precisely, 10 μg of each DNA was mixed with pure distilled water, making the final volume 225 μl . Then, 25 μl of 2.5 M CaCl_2 solution was added and mixed by vortexer. 250 μl of 2x HBS solution (280 mM NaCl, 10 mM KCl, 1.5 mM $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 12 mM dextrose, 50 mM HEPES) was slowly dropped drop by drop into a tube containing the DNA. Reaction was induced for 10 minutes at room temperature. The mixture was transferred onto a cell culture dish, followed by further culture for 6 hours. The dish was

washed with PBS (phosphate buffered saline) three times and then a fresh medium was supplemented, followed by further culture for 48 more hours.

5 <Example 3> Confirmation of hFSH expression by ELISA

Temporarily over-expressed FSH was quantified by using the culture medium of COS-7 cells transfected with Rc/CMV-dhfr-TPL-hFSH beta/alpha plasmid or Rc/CMV-dhfr-TPL (control plasmid) in the above example 2.

10 ELISA (enzyme-linked immunosorbent assay) kit (MEDI-CORP, cat# KTSF 3651, Canada) was used to detect FSH, by following the manufacturer's protocol. Particularly, 100 μ l of the transfected cell culture medium cultured for 48 hours and 100 μ l of standard solution equipped

15 in the kit were added to the well coated with anti-human FSH, leading to the reaction for 30 minutes on a plate shaker. The wells were washed with washing solution provided by the manufacturer, and then, anti-human FSH antibody linked to HRP (horse radish

20 peroxidase) was added to each well by 100 μ l, followed by reaction on the plate shaker at room temperature. After 30 minutes of reaction, the plate was washed with washing solution three times. Then, 100 μ l of substrate solution was added to each well and reaction was

25 induced at room temperature on the plate shaker. 15

minutes later, 100 μ l of stop solution was added to each well and OD₄₅₀ was measured. Using the standard solution provided by the manufacturer of the kit, standard curve and function were obtained. The amount of FSH was quantified by using the standard curve and the function.

As a result, human FSH was hardly shown in cells transfected with Rc/CMV-dhfr-TPL, a control plasmid, but human FSH was over-expressed in cells transfected with Rc/CMV-dhfr-TPL-hFSH beta/alpha plasmid (FIG. 4).

<Example 4> Preparation of a CHO cell line expressing hFSH

Rc/CMV-dhfr-TPL-hFSH beta/alpha vector, which was proved in the above example 3 to over-express hFSH, was introduced into Chinese hamster ovary cells (CHO/dhfr⁻) (Urlaub et al., *Somat. Cell Mol. Genet.*, 12:555-566, 1986) having incomplete nucleic acid biosynthesizing process resulted from the damaged DHFR gene, leading to the preparation of a cell line suitable for mass-production of hFSH protein. More precisely, before a hFSH over-expressing vector was introduced, CHO/dhfr⁻ cells were thawed and put in a T-75 culture flask, followed by culture in MEM-alpha medium (GIBCO BRL,

USA) supplemented with HT supplement (GIBCO BRL, USA) and 10% dialyzed fetal bovine serum (FBS) until cell density reached 80-90%. Trypsin was treated to those cells growing with being adhered to the plate. Then, cells were recovered and 5×10^5 cells were plated on 60 mm dish. The cells were transfected with Rc/CMV-dhfr-TpL-hFSH beta/alpha, a FSH expression vector, by means of CaPO_4 coprecipitation explained in the above example 3, followed by further culture for two more days. After the completion of the culture, the transfected CHO/dhfr⁻ cells were diluted consecutively (1/2, 1/5, 1/10, 1/20, 1/50, 1/100, 1/200, 1/500), and then, cultured in HT supplement-free medium on 100 mm dish, during which the primary screening was performed. During screening, the medium was replaced once a week, passage was omitted, and plates with colony formed were selected after 2 to 3 weeks. Colonies were separated from the plates and transferred into a 24-well plate for subculture. About one or two weeks later, cell clones in the 24-well plate were treated with trypsin. Then, cells were recovered and plated again in a 24-well plate (2×10^4 cells/well). 48 hours later, supernatant was obtained to measure the expression level of FSH by using FSH detecting ELISA kit (MEDI-CORP, cat# KTSF 3651, Canada). Cell clones showing

higher level of FSH expression were diluted consecutively, which were put in a 100 mm dish for further culture. 20 nM of methotrexate (MTX) was added to the medium for the second screening. Finally,
5 clones high-expressing human FSH were selected with the same identifying processes as used for the primary screening.

<Example 5> Selection of a CHO cell line expressing

10 hFSH high

CHO/dhfr⁻ cells deficient in DHFR cannot survive in MEM-alpha medium, a minimum essential medium devoid of nucleic acid, because the cells have an incomplete nucleic acid biosynthesizing process. However,
15 recombinant CHO cells in which FSH gene and DHFR gene are inserted, recover nucleic acid biosynthesizing capability, so that the MEM-alpha medium can be used for the selection culture. Thus, in the present invention, in order to select cell lines expressing
20 hFSH high, clones prepared in the above example 4 were cultured in culture media having different MTX concentrations (20 nM to 1 μ M). More precisely, the primary selected clones were subcultured in MEM-alpha medium supplemented with 10% of dialyzed FBS, during

which the concentration of MTX was increased gradually. Clones were cultured at different MTX concentrations to select specific ones having resistance against MTX. The supernatants of each clone in different culture stages were obtained to measure the expression of FSH, resulting in the second selection of cells expressing FSH high. The expression of FSH was quantified by means of ELISA described in the above example 3.

For the third selection, monoclonal selection was performed by means of limiting dilution method to reduce the chance of heterogeneity during MTX amplification. That is, single cells having even FSH expression rate and excellent productivity were separated from all the clones showing irregular FSH expression rates by means of limiting dilution method. Cells were distributed in a 96-well plate by one-two clones per each well, and then the wells inoculated with 1 cell only were selected, followed by further culture for 2 to 3 weeks. From the plate showing colonies formed, cells were separated and sub-cultured to measure the amount of expressed FSH by ELISA. At last, single cells highly expressing FSH were separated (FIG. 5). Among those separated cells through the third selection, a recombinant CHO cell line showing a stable proliferation in the presence of 1 μ M MTX and

high FSH productivity was selected and named "DPFC325", which was deposited at Korean Cell Line Research Foundation (KCLRF), Seoul, Korea on June 20, 2003 (Accession No : KCLRF-BP-00082 ㉔).

5

<Example 6> Confirmation of the insertion of hFSH gene
in chromosome by FISH

For the clones used for the production of a protein as a treating agent, genetic stability is so
10 important. In order for a plasmid vector inserted in a cell line to be transmitted to the next generation stably, integration in chromosome of CHO cell is essential. In the present invention, FISH (fluorescent *in situ* hybridization) was performed to investigate
15 whether FSH gene was inserted in chromosome of a recombinant cell line or not. Precisely, metaphase chromosome of CHO cell and nucleus of interphase were hybridized with biotin-labeled probe. The hybridized metaphase chromosome was reacted with avidin conjugated
20 with a fluorescent material, followed by observation.

As a result, almost all nuclei of interphase showed reaction color, indicating that FSH gene was successfully inserted in chromosome (FIG. 6).

Industrial Applicability

As explained hereinbefore, a transformant of the present invention can be effectively used for the treatment of infertility since the transformant can
5 produce human follicle stimulating hormone stably and largely.

Sequence List Text

Nucleotide sequences represented by SEQ. ID. No 1
10 to SEQ. ID. No 15 are the ones used for the construction of recombinant hFSH expression vector (Rc/CMV-dhfr-TPL-hFSH beta/alpha) of the example 1.

Those skilled in the art will appreciate that the
15 conceptions and specific embodiments disclosed in the foregoing description may be readily utilized as a basis for modifying or designing other embodiments for carrying out the same purposes of the present invention. Those skilled in the art will also appreciate that such
20 equivalent embodiments do not depart from the spirit and scope of the invention as set forth in the appended claims.


Identification of Deposit

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEPTION IN THE CASE OF AN ORIGINAL DEPOSIT
issued pursuant to Rule 7.1

To: ProGen Co Ltd.
47-5 Sanggal-Ri, Kiheung-Up.
Yongin-Si, Kyunggi-Do, 449-900 KOREA

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : DPFC325	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCLRF-BP-00082
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by : <input type="checkbox"/> A scientific description <input type="checkbox"/> A proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on June 20, 2003.	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Director Korean Cell Line Research Foundation Address : Cancer Research Institute Seoul National University College of Medicine 28 Yongon-dong, Chongno-Gu Seoul, 110-744, Korea	Signature(s) :  Date : 2003. 6. 23.

Form BP/4 (KCLRF Form 17)

Page sole

What is claimed is

1. An expression vector to express human follicle stimulating hormone (FSH) comprising a gene coding
5 human FSH, a promoter sequence, a polyadenylation motif sequence and a dihydrofolate reductase (DHFR) gene.

2. The expression vector as set forth in claim 1, wherein the human FSH gene consists of human FSH alpha
10 subunit gene represented by SEQ. ID. No 1, internal ribosomal entry site (IRES) sequence represented by SEQ. ID. No 7 and human FSH beta subunit gene represented by SEQ. ID. No 2.

15 3. The expression vector as set forth in claim 1, wherein the promoter is the promoter of early gene of cytomegalovirus (CMV) represented by SEQ. ID. No 8.

4. The expression vector as set forth in claim 1,
20 wherein tripartite leader sequence of adenovirus represented by SEQ. ID. No 9 is additionally included.

5. The expression vector as set forth in claim 1, wherein the polyadenylation motif is a polyadenylation
25 motif of early gene of SV40 virus, represented by SEQ.

ID. No 13, and/or a polyadenylation motif of bovine growth hormone (BGH) gene, represented by SEQ. ID. No 14.

5 6. The expression vector as set forth in claim 1, wherein the dihydrofolate reductase (DHFR) gene is a base sequence represented by SEQ. ID. No 12.

 7. The expression vector as set forth in claim 1,
10 wherein the vector is RC/CMV-dhfr-TPL-hFSH beta/alpha represented in FIG. 1.

 8. A recombinant transformant mass-producing human FSH prepared by introducing the expression vector
15 of claim 1 into host cells.

 9. The recombinant transformant as set forth in claim 8, wherein the host cell is a CHO (Chinese hamster ovary) originated cell line (CHO/dhfr⁻)
20 harboring damaged dihydrofolate reductase (DHFR) gene.

 10. A recombinant transformant DPFC325 (Accession No: KCLRF-BP-00082) mass-producing human FSH prepared by introducing the expression vector of claim 7 into a
25 CHO originated cell line (CHO/dhfr⁻).

11. A method for mass-production of human follicle stimulating hormone comprising the following steps:

- 5 1) Preparation of an expression vector containing human FSH gene;
- 2) Transfection of host cells with the expression vector of the step 1);
- 3) Selection of transformants transfected in the
10 step 2);
- 4) Selection of a recombinant transformant stably producing human FSH from the recombinant transformants selected in the step 3); and
- 5) Obtainment of human FSH from the culture of
15 the recombinant transformant selected in the step 4).


12. The method for mass-production of human follicle stimulating hormone as set forth in claim 11, wherein the expression vector of step 1) is an
20 expression vector of claim 1.

13. The method for mass-production of human follicle stimulating hormone as set forth in claim 11, wherein the host cell of step 2) is a CHO originated

cell line (CHO/dhfr⁻) harboring damaged dihydrofolate reductase (DHFR) gene.

14. The method for mass-production of human
5 follicle stimulating hormone as set forth in claim 11,
wherein the selected recombinant transformant of step
4) is a transformant of anyone of claim 8 to 10.

1/6



2/6

Wild type DNA sequence	ATG GAT TAC TAC AGA AAA TAT GCA GCT ATC TTT CTG GTC ACA TTG
Amplified DNA sequence	---
Amino acid sequence	Met Asp Tyr Tyr Arg Lys Tyr Ala Ala Ile Phe Leu Val Thr Leu
	TCG GTC TTT CTG CAT GTT CTC CAT TCC GCT CCT GAT GTG CAG CAT

	Ser Val Phe Leu His Val Leu His Ser Ala Pro Asp Val Gln Asp
	TGC CCA GAA TGC ACG CTA CAG GAA AAC CCA TTC TTC TCC CAG CCG

	Cys Pro Glu Cys Thr Leu Gln Glu Asn Pro Phe Phe Ser Gln Pro
	GGT GCC CCA ATA CTT CAG TGC ATG GGC TGC TGC TTC TCT AGA GCA

FSH α subunit	Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys Phe Ser Arg Ala
	TAT CCC ACT CCA CTA AGG TCC AAG AAG ACG ATG TTG GTC CAA AAG

	Tyr Pro Thr Pro Leu Arg Ser Lys Lys Thr Met Leu Val Gln Lys
	AAC GTC ACC TCA GAG TCC ACT TGC TGT GTA GCT AAA TCA TAT AAC

	Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser Tyr Asn
	AGG GTC ACA GTA ATG GGG GGT TTC AAA GTG GAG AAC CAC ACG GCG


	Arg Val Thr Val Met Gly Gly Phe Lys Val Glu Asn His Thr Ala
	TGC CAC TGC AGT ACT TGT TAT TAT CAC AAA TCT TAA

	Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser termination


3/6

Wild type DNA sequence	ATG AAG ACA CTC CAG TTT TTC TTC CTT TTC TGT TGC TGG AAA GCA
Amplified DNA sequence	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
Amino acid sequence	Met Lys Thr Leu Gln Phe Phe Phe Leu Phe Cys Cys Trp Lys Ala
	ATC TGC TGC AAT AGC TGT GAG CTG ACC AAC ATC ACC ATT GCA ATA
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Ile Cys Cys Asn Ser Cys Glu Leu Thr Asn Ile Thr Ile Ala Ile
	GAG AAA GAA GAA TGT CGT TTC TGC ATA AGC ATC AAC ACC ACT TGG
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Glu Lys Glu Glu Cys Arg Phe Cys Ile Ser Ile Asn Thr Thr Trp
	TGT GCT GGC TAC TGC TAC ACC AGG GAT CTG GTG TAT AAG GAC CCA
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Cys Ala Gly Tyr Cys Tyr Thr Arg Asp Leu Val Tyr Lys Asp Pro
FSH β subunit	GCC AGG CCC AAA ATC CAG AAA ACA TGT ACC TTC AAG GAA CTG GTA
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Ala Arg Pro Lys Ile Gln Lys Thr Cys Thr Phe Lys Glu Leu Val
	TAT GAA ACA GTG AGA GTG CCC GGC TGT GCT CAC CAT GCA GAT TCC
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Tyr Glu Thr Val Arg Val Pro Gly Cys Ala His His Ala Asp Ser
	TTG TAT ACA TAC CCA GTG GCC ACC CAG GTG CAC TGT GGC AAG TGT
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Leu Tyr Thr Tyr Pro Val Ala Thr Gln Val His Cys Gly Lys Cys
	GAC AGC GAC AGC ACT GAT TGT ACT GTG CGA GGC CTG GGG CCC AGC
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Asp Ser Asp Ser Thr Asp Cys Thr Val Arg Gly Leu Gly Pro Ser
	TAC TGC TCC TTT GGT GAA ATG AAA GAA TAA
	--- --- --- --- --- --- --- --- --- --- --- --- --- --- ---
	Tyr Cys Ser Phe Gly Glu Met Lys Glu termination


4 / 6



5/6



6/6



Sequence Listing

<110> ProGen Co Ltd.

5 <120> Method for mass production of human Follicle Stimulating Hormone

<150> KR 10-2003-0068641

<151> 2003-10-02

10 <160> 15

<170> KopatentIn 1.71

<210> 1

15 <211> 351

<212> DNA

<213> Homo sapiens

<400> 1

20 atggattact acagaaaata tgcagctatc ttctctgggtca cattgtcggg ctttctgcat 60

gtttctccatt ccgctcctga tgtgcaggat tgcccagaat gcacgctaca ggaaaaccca 120

ttctttctccc agccgggtgc cccaatactt cagtgcattgg gctgctgctt ctctagagca 180

25

tatccactc cactaaggtc caagaagacg atgttgggtcc aaaagaacgt cacctcagag 240
 tccacttgct gtgtagctaa atcatataac aggggtcacag taatggggggg ttcaaagtg 300
 5 gagaaccaca cggcgtgcc a ctgcagtact tgttattatc acaaattcta a 351
 .
 <210> 2
 <211> 390
 10 <212> DNA
 <213> Homo sapiens
 .
 <400> 2
 atgaagacac tccagttttt ctcccttttc tgttgctgga aagcaatctg ctgcaatagc 60
 15 tgtgagctga ccaacatcac cattgcaata gagaaagaag aatgtcgitt ctgcataagc 120
 atcaacacca ctgtggtgtgc tggctactgc tacaccaggg atctggtgta taaggacca 180
 20 gccaggccca aaatccagaa aacatgtacc ttcaaggaac tggatatga aacagtgaga 240
 gtgcccggt gtgctacca tgcagattcc ttgtatacat acccagtggc caccaggtg 300
 cactgtggca agtgtgacag cgacagcact gattgtactg tgcgaggcct gggggccagc 360
 25

tactgctcct ttggtgaaat gaaagaataa

390

<210> 3

5 <211> 27

<212> DNA

<213> Artificial Sequence

<220>

10 <223> sense primer for human FSH alpha subunit

<400> 3

agcgccatgg attactacag aaaatat

27

15

<210> 4

<211> 37

<212> DNA

20 <213> Artificial Sequence

<220>

<223> antisense primer for human FSH alpha subunit

25

<400> 4
gggggatccg gggcccttaag attgtgata attaaca 37

5 <210> 5
<211> 33
<212> DNA
<213> Artificial Sequence

10 <220>
<223> sense primer for human FSH beta subunit

<400> 5
15 gggggcgccc gcaggatgaa gacactccag ttt 33

<210> 6
<211> 27
20 <212> DNA
<213> Artificial Sequence

<220>
<223> antisense primer for human FSH beta subunit
25

	<400>	6	
	ggggataatct tattctttca ttccacc		27
5			
	<210>	7	
	<211>	592	
	<212>	DNA	
	<213>	Encephalomyocarditis virus	
10			
	<400>	7	
	gatatcgaat tccccctctc cctcccccc ccctaacgtt actggcogaa gccgcttga		60
	ataaggccgg tgtgcgtttg tctat atgtt atttccacc atattgccgt cttttggcaa		120
15			
	tgtgagggcc cggaaacctg gccctgtctt ctgacgagc attcctaggg gtctttcccc		180
	tctcgccaaa ggaatgcaag gtctgttgaa tgcgtgaag gaagcagttc ctctggaagc		240
20			
	ttcttgaaga caaacaacgt ctgtagcgac cctttgcagg cagcgggaacc cccacctgg		300
	cgacagggtgc ctctcgggcc aaaagccacg tgtataagat acacctgcaa aggcggcaca		360
	acccagtgac caggttgtga gttggatagt tgtggaaaga gtcaaatggc tctcctcaag		420
25			

cgtattcaac aaggggctga aggatgcca gaaggtagcc cattgtatgg gatctgatct 480

ggggcctcgg tgcacatgct ttacatgtgt ttagtcgagg ttaaaaaaac gtctaggccc 540

5 cccgaaccac ggggacgtgg ttttcctttg aaaaacacga tgataaatgg cc 592

<210> 8

<211> 654

10 <212> DNA

<213> Cytomegalovirus

<220>

<221> promoter

15 <222> (1)..(654)

<400> 8

gatgtacggg ccagatatag gcgttgacat tgattatga ctagttatta atagtaatca 60

20 attacggggc cattagtcca tagcccatat atggagtacc gcgttacata acttacggta 120

aatggccgc ctggctgacc gcccaacgac ccccgcccat tgacgtcaat aatgacgtat 180

gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga ctatttacgg 240

25

	taaactgccc acttggcagt acatcaagtg tatcatatgc caagtacgcc ccctattgac	300
	gtcaatgacg gtaaattggcc cgcttgcat tatgccagc acatgacctt atgggacttt	360
5	cctacttggc agtacatcta cgtattagtc atcgctatta ccatggatgat gcggttttgg	420
	cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag tctccacccc	480
	attgacgtca atgggagttt gttttggcac caaatcaac gggactttcc aaaatgtcgt	540
10	aacaactccg ccccatgac gcaaatgggc ggtaggcgtg tacgggtggga ggtctatata	600
	agcagagctc tctggctaac tagagaaccc actgcttaac tgcttatoga aatt	654
15	<210> 9	
	<211> 441	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Adenovirus tripartite leader sequence	
25	<400> 9	

togatactct ctccgcac gcgtctgcg agggccagct gttgggctcg cggttgagga 60
 caaactcttc gcggtcttc cagtactctt ggalcggaaa ccgctcgcc tccgaacggt 120
 5 actccgccac cgaggacct gagcgagtc gcatcgaccg gatcggaaaa cctctcgact 180
 gtgggggtga gtactccctc taaaagcgg gcatgacttc tgcgctaaga ttgtcagttt 240
 ccaaaaacga ggaggatttg atattcacct gggccgcggt gatgcctttg agggtgccg 300
 10 cgtccatctg gtcagaaaag acaatctttt tttgtcaag cttgagggtg ggcaggcttg 360
 agatctggcc atacactga gtgacaatga catccacttt gcctttctct ccacagggtg 420
 15 ccactcccag gtccaactgc a 441

<210> 10
 <211> 19
 20 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> sense primer for tripartite leader sequence
 25

<400> 10
 gatatcgata ctctcttcc 19

 5
 <210> 11
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> antisense primer for tripartite leader sequence

 15 <400> 11
 gcgtcgacct gcagttggac ctgggag 27

 <210> 12
 20 <211> 564
 <212> DNA
 <213> Artificial Sequence

 <220>
 25 <223> dihydrofolate reductase

<400> 12

atgggtcgcac cattgaactg catcgctgcc gtgtcccaaa atatgggat tggcaagaac 60

5 ggagacctac cctggcctcc gtcaggaac gagtcaagt acttccaaag aatgaccaca 120

acctcttcag tggaaggtaa acagaatctg gtgattatgg gtaggaaaac ctggttctcc 180

10 attcctgaga agaatcgacc tttaaaggac agaattaata tagttctcag tagagaactc 240

aaagaaccac cacgaggagc tcattttctt gccaaaagt tggatgatgc cttaagactt 300

attgaacaac cggaattggc aagtaaagta gacatggtit ggatagtcgg aggcagttct 360

15 gtttaccagg aagccatgaa tcaaccaggc caccctcagac tctttgtgac aaggatcatg 420

caggaatttg aaagtgcac gtttttccca gaaattgatt tggggaaata taaacttctc 480

20 ccagaatacc caggcgtcct ctctgaggtc caggaggaaa aaggcatcaa gtataagttt 540

gaagtctacg agaagaaaga cttaa 564

25 <210> 13

<211> 130
<212> DNA
<213> Simian virus 40

5 <220>
<221> polyA_signal
<222> (1)..(130)

<400> 13

10 aacttgttta ttgcagctta taatggttac aaataaagca atagcatcac aaatttcaca 60
aataaagcat ttttttcaat gcattctagt tgtggtttgt ccaaactcat caatgtatct 120
tatcatgtct 130

15

<210> 14
<211> 232
<212> DNA
20 <213> Artificial Sequence

<220>
<223> bovine growth hormone

25

<220>
 <221> polyA_signal
 <222> (1)..(232)

5 <400> 14
 ctagagctcg ctgacagcc tcgactgtgc cttctagttg ccagccatct gttgtttgcc 60
 cctcccccgt gccttccttg accctggaag gtgccactcc cactgtcctt tcctaataaa 120
 -
 10 atgaggaaat tgcacgcat tgtctgagta ggtgtcattc tattctgggg ggtgggggtg 180
 ggcaggacag caagggggag gattgggaag acaatagcag gcatgctggg ga 232

15 <210> 15
 <211> 10
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> BamH I linker

 <400> 15
 25 cgggatcccg 10

INTERNATIONAL SEARCH REPORT

International application No.

PCT/KR2004/002474

A. CLASSIFICATION OF SUBJECT MATTER**IPC7 C12N 15/85**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC : C12N, C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean Patents and Applications for Inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PubMed, Delphion, CA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Gebert, A., et al. "Expression of FSH in CHO cells, I. Comparison of promoter types and effects of their respective inducers" Cytotechnology, 1994, Vol. 14(1): pages 39-45, See entire document.	1-14
A	Mannaerts, B., et al. "Comparative in vitro and in vivo studies on the biological characteristics of recombinant human follicle-stimulating hormone" Endocrinology, 1991, Vol. 129(5): pages 2623-2630, See entire document	1-14
A	US 2001/0007757A1 (Boime et al.), Jul. 12, 2001, See entire document	1-14

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 JANUARY 2005 (13.01.2005)

Date of mailing of the international search report

13 JANUARY 2005 (13.01.2005)

Name and mailing address of the ISA/KR

Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

PARK, JEONG UNG

Telephone No. 82-42-481-8159



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR2004/002474

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2001/0007757A1	Jul. 12, 2001	US 6306654	Oct. 23, 2001